

1. Let  $k=5$ . Assume a cache with initial contents 1,2,3,4,5 and with at least one item not in the cache. Give an example sequence of length at least 6 of cache requests and a sequence of random choices by the Marker algorithm so that its competitiveness on this sequence with these choices would be greater than H5.

Q1

$k=5$

cache has initial contents 

1	2	3	4	5
---	---	---	---	---

let there  $k+1$  length = 6.

let the request sequence be.

2, 3, 2, 5, 6, 4, 1, 4, 6, 3, 2, 5, 1

Marker Algorithm

	2	3	2	5	6	4		1	4	6	3	2		5	1
0	0	0	0	0	1	1	R	0	1	1	1	1	1	R	0
0	1	1	1	1	1	1	E	0	0	0	1	1	1	E	0
0	0	1	1	1	1	1	S	0	0	0	0	1	1	S	0
0	0	0	0	0	0	1	E	0	0	1	1	1	1	E	0
0	0	0	0	1	1	1	T	0	0	0	0	0	1	T	0

1
2
3
4
5

6
2
3
4
5

1
2
3
4
5

1
6
3
4
5

1
6
3
4
2

5
6
3
4
2

5
1
3
4
2

initial  
cache

- evict the lowest indexed cache
- when all the cache pages are marked it resets all to unmarked
- there are 6 cache misses

MIN algorithm

cache 

1	2	3	4	5
---	---	---	---	---

✓ → no cache miss

x → cache miss, evict the item farthest in future

2	3	2	5	6	4	1	4	6	3	2	5	1
✓	✓	✓	✓	x	✓	✓	✓	✓	✓	✓	x	✓

1
2
3
4
6

evict 5

1
5
3
4
6

evict 2

- number of misses is 2

The length of the request sequence  $N = 13$

MIN can have  $\frac{N}{K}$  misses in the worst case

$$\frac{N}{K} = \frac{13}{5} = 2.6$$

For our request sequence MIN faults 2 times.

For the same request sequence Marker Algorithm faults 6 times

- We call an item stale if it is unmarked, but was marked in the previous round and clean if its neither marked nor is it stale.
- Let  $m$  be the number of requests to clean items in a round.

$$C_{H_5} = 2 \cdot \lceil \text{MIN} \rceil \quad C_{MA} = 6 \cdot [\text{Marker Algorithm}]$$

$\therefore$  competitiveness of Marker Algorithm is greater than  $H_5$ .

$$C_{H_5} < C_{MA}$$

$$\text{or } \boxed{C_{MA} > C_{H_5}}$$

2. Use the extended Euclidean algorithm to find the multiplicative inverse of  $26 \bmod 1155$ .  
Solve  $9x \equiv 6 \bmod 33$  for all solutions.

**Extended-Euclid(a,b)**

1. if  $b = 0$  then return  $(a, 1, 0)$
2.  $(d', x', y') = \text{Extended-Euclid}(b, a \bmod b)$
3.  $(d, x, y) = (d', y', x' - \text{floor}(a/b) * y')$
4. return  $(d, x, y)$

Q2

Part 1

EE(26, 1155)

$(d, x', y') = \text{EE}(1155, 26)$

$(d, x', y') = \text{EE}(26, 11)$

$(d, x', y') = \text{EE}(11, 4)$

$(d, x', y') = \text{EE}(4, 3)$

$(d, x', y') = \text{EE}(3, 1)$

$(d, x', y') = \text{EE}(1, 0)$

$(d, x, y) = (1, 1, 0)$

$(d, x, y) = (1, 0, 1)$

$(d, x, y) = (1, 1, -1)$

$(d, x, y) = (1, -1, 3)$

$(d, x, y) = (1, 3, -7)$

$(d, x, y) = (1, -7, 311)$

$(d, x, y) = (1, 311, -7)$

EE(26, 1155) =  $(1, 311, -7)$

where  $d = 1$   $x = 311$   $y = -7$

$$26 \times 311 - 1155 \times 7$$

$$8086 - 8085 = 1 \quad [ax + by = d]$$

$$\boxed{ax = 1 \bmod b}$$

Here  $x$  is the multiplicative inverse of  $26 \bmod 1155$

$\boxed{x = 311}$  is multiplicative inverse of  $26 \bmod 1155$

## Part 2

Solve  $ax \equiv b \pmod{n}$   
 $ax = b \pmod{n}$

EE(9, 33)

$$(d, x', y') = \text{EE}(33, 9)$$

$$(d, x', y') = \text{E}(9, 6)$$

$$(d, x', y') = \text{EE}(6, 3)$$

$$(d, x', y') = \text{EE}(3, 0)$$

$$(d, x, y) = (3, 1, 0)$$

$$(d, x, y) = (3, 0, 1)$$

$$(d, x, y) = (3, 1, -1)$$

$$(d, x, y) = (3, -1, 4)$$

$$(d, x, y) = (3, 4, -1)$$

Now executing Modular-Linear-Equation-Solver (a, b, n)  
for  $ax \equiv b \pmod{n}$

$$a = 9 \quad b = 6 \quad n = 33$$

$$x_0 = x(b/d) \pmod{n}$$

$$= 4(6/3) \pmod{33}$$

$$x_0 = 8$$

Now executing the for loop  $i = 0$  to  $d-1$   
 $i = 0$  to  $2$

$$\text{on } x_i = x_0 + i * (b/d) \pmod{n}$$

$$i = 0 \quad x_0 = 8$$

$$i = 1 \quad x_1 = (8 + 1 * 33/3) \pmod{33} = (8 + 11) \pmod{33}$$

$$x_1 = 19$$

$$i = 2 \quad x_2 = (8 + 2 * \frac{33}{3}) \pmod{33} = (8 + 22) \pmod{33}$$

$$x_2 = 30$$

so the solutions are 8, 19, 30

3. Using the Chinese Remainder theorem, determine a number  $x \bmod 1155$  that satisfies  $x \equiv 2 \bmod 3$ ,  $x \equiv 3 \bmod 5$ , and  $x \equiv 4 \bmod 7$ , and  $x \equiv 5 \bmod 11$ .

Q3

$x \bmod 1155$  that satisfies

$$x \equiv 2 \bmod 3$$

$$x \equiv 3 \bmod 5$$

$$x \equiv 4 \bmod 7$$

$$x \equiv 5 \bmod 11$$

Calculate  $m_i$

$$m_1 = \frac{1155}{3} = 385$$

$$m_2 = \frac{1155}{5} = 231$$

$$m_3 = \frac{1155}{7} = 165$$

$$m_4 = \frac{1155}{11} = 105$$

Calculate  $t_i$

$$t_i = (m_i)^{-1} \bmod n_i$$

For  $i=1$

EE(385, 3)

$$(d, x', y') = \text{EE}(3, 1)$$

$$(d, x', y') = \text{EE}(1, 0)$$

$$(d, x, y) = (1, 1, 0)$$

$$(d, x, y) = (1, 0, 1)$$

$$(d, x, y) = (1, 1, -128)$$

This tells us  $385 \times 1 - 3 \times 128 = 1$   
 $\equiv 385 \times 1 = 1 \bmod 3$

$$t_1 = 1 \bmod 3$$

For  $t_i$  when  $i = 2$

$$EE(231, 5) = (1, 1, -46)$$

$$d=1, \quad x=1, \quad y=-46$$

$$\begin{aligned} \text{which gives us } 231 \times 1 - 5 \times 46 &= 1 \\ &\equiv 231 \times 1 = 1 \pmod{5} \end{aligned}$$

$$t_2 = 1 \pmod{5}$$

when  $i = 3$

$$EE(165, 7) = (1, 2, -47)$$

$$d=1 \quad x=2 \quad y=-47$$

$$\begin{aligned} \text{which gives us } 165 \times 2 - 7 \times 47 &= 1 \\ &\equiv 165 \times 2 = 2 \pmod{7} \end{aligned}$$

$$t_3 = 2 \pmod{7}$$

when  $i = 4$

$$EE(105, 11) = (1, 2, -19)$$

$$\begin{aligned} \text{We get } 105 \times 2 - 11 \times 19 &= 1 \\ &\equiv 105 \times 2 = 2 \pmod{11} \end{aligned}$$

$$t_4 = 2 \pmod{11}$$

Calculate  $c_i$

$$\begin{aligned} c_1 &= m_1 t_1 \\ &= 385 \times 1 \\ &= 385 \end{aligned}$$

$$\begin{aligned} c_2 &= m_2 t_2 \\ &= 231 \times 1 \\ &= 231 \end{aligned}$$

$$\begin{aligned} c_3 &= m_3 t_3 \\ &= 165 \times 2 \\ &= 330 \end{aligned}$$

$$\begin{aligned} c_4 &= m_4 t_4 \\ &= 105 \times 2 \\ &= 210 \end{aligned}$$

$$\begin{aligned} x &= a_1 c_1 + a_2 c_2 + a_3 c_3 + a_4 c_4 \\ &= (2 \times 385) + (3 \times 231) + (4 \times 330) + (5 \times 210) \\ &= 770 + 693 + 1320 + 1050 \\ &= 3833 \end{aligned}$$

Now substituting  $x$  in  $x \bmod 1155$

we get  $\boxed{3833 \bmod 1155 = 365}$

$$x \bmod 3 = 368 \bmod 3 \equiv 2 \bmod 3$$

---

$$x \bmod 5 = 368 \bmod 5 \equiv 3 \bmod 5$$

$$x \bmod 7 = 368 \bmod 7 \equiv 4 \bmod 7$$

$$x \bmod 11 = 368 \bmod 11 \equiv 5 \bmod 11$$

4. Suppose  $p=7, q=17$ . If we choose  $e=3$ , what would be the RSA public and private keys? Show the result of encrypting with the private key, the message 89. Show the steps in decrypting it, to get the original number back.

1. Select two large prime numbers  $p$  and  $q$  such that  $p \neq q$ . (For instance, the primes might be 512 bits each.)
2. Compute  $n = pq$ .
3. Select a small odd integer  $e$  that is relatively prime to  $\phi(n) = (p-1)(q-1)$ .
4. Compute the multiplicative inverse  $d$  of  $e \bmod \phi(n)$ .
5. Publish the pair  $P = (e, n)$  as the RSA public key.
6. Keep secret the pair  $S = (d, n)$  as the RSA secret key.

Q4

step 1:  $p$  and  $q$  are both primes and  $p \neq q$

step 2:  $n = p * q$   
 $= 7 * 17 = 119$

step 3:  $\phi(n) = (p-1)(q-1) = 6 * 16 = 96$ , we have  $e=5$   
 $\therefore \phi(n), e$  are relatively prime.  
 $\gcd(5, 96) = 1$

step 4 compute multiplicative inverse  $d$  of  $e \bmod \phi(n)$   
 when  $\gcd(a, n) = 1$  (Special case) we have  $b=1$ .  
 $\therefore ax \equiv b \pmod{n}$  [ $a=5, b=1, n=96$ ]  
 we have.

$$5x \equiv 1 \pmod{96} \rightarrow (\text{eq 1})$$

We solve the above using Modular Linear equation solver  
 MLES ( $a, b, n$ )

EE( $a, n$ )

EE( $5, 96$ )

$$(d, x', y') = \text{EE}(96, 5)$$

$$(d, x', y') = \text{EE}(5, 1)$$

$$(d, x', y') = \text{EE}(1, 0)$$

$$(d, x, y) = (1, 1, 0)$$

$$(d, x, y) = (1, 0, 1)$$

$$(d, x, y) = (1, 1, -19)$$

$$(d, x, y) = (1, -19, 1)$$

$$x_0 = (x \cdot (b/a)) \bmod n$$

$$= -19 \bmod 96$$

$$= 77$$

$$\therefore \boxed{d = 77}$$



Step 5: public key  $P = (e, n) = (5, 119)$

Step 6: Secret key  $S = (d, n) = (77, 119)$

Encryption of Message  $M=89$  with private key.

To apply a key to a message  $0 \leq M < n$ , we

compute either  $P(M) = M^e \bmod n$  or  $S(C) = C^d \bmod n$ .

For our encryption we use  $P(M) = M^e \bmod n$

Here  $M=89$   $e=5$   $n=119$

substituting the values in the above equation

$$\begin{aligned} P(M) &= 89^5 \bmod 119 \\ &= 5584059449 \bmod 119 \\ &= 38 \end{aligned}$$

Decryption steps to get  $M=89$

we solve this using Exp-Mod( $a, x, n$ ) algorithm.

$$38^{77} \bmod 119$$

$$R1: \text{Exp-Mod}(38, 77, 119)$$

$$C = (\text{Exp-Mod}(38, 38, 119))^2 \bmod 119$$

$$R2: \text{Exp-Mod}(38, 38, 119)$$

$$C = (\text{Exp-Mod}(38, 19, 119))^2 \bmod 119$$

$$R3: \text{Exp-Mod}(38, 19, 119)$$

$$C = (\text{Exp-Mod}(38, 9, 119))^2 \bmod 119$$

$$R4: \text{Exp-Mod}(38, 9, 119)$$

$$C = (\text{Exp-Mod}(38, 4, 119))^2 \bmod 119$$

$$R5: \text{Exp-Mod}(38, 4, 119)$$

$$C = (\text{Exp-Mod}(38, 2, 119))^2 \bmod 119$$

R6: Exp-Mod (38, 2, 119)

$$C = (\text{Exp-Mod}(38, 1, 119))^2 \bmod 119$$

R7: Exp-Mod (38, 1, 119)

$$x = 1 \text{ so return } 38 \bmod 119 = 38$$

Recurising back to R4

$$R6: C = 38^2 \bmod 119 = 16$$

$x$  is even so return  $C$

$$R5: C = 16^2 \bmod 119 = 18$$

$x$  is even so return  $C$

$$R4: C = 18^2 \bmod 119 = 86$$

$x$  is odd so return  $a * C \bmod n$

$$38 * 86 \bmod 119 = 55$$

return 55

$$R3: C = 55^2 \bmod 119 = 50$$

$x$  is odd. so return  $(38 * 50) \bmod 119 = 115$

$$R2: C = 115^2 \bmod 119 = 16$$

$x$  is even so return  $C$

$$R1: 16^2 \bmod 119 = 18$$


$x$  is odd so return  $(38 * 18) \bmod 119 = \underline{\underline{89}}$

So the message is decrypted and we get back.

$$\boxed{M=89}$$

Coding Question:

By Sylvester's Theorem, we know there is always a prime in this range. For 0.5pts of your coding points described below, plot  $\log_2(x-2^{n-1})$  vs  $n$  where  $x$  is your output for values  $n$  between 100 and 200. Say what you observe.



From the above graph we can say that  $\log_2(x-2^{n-1})$  linearly increases as  $n$  increases. So its  $O(n)$ .