

On the edge-magic indices of $(v, v+1)$ -graphs

Sin-Min Lee

Department of Computer Science
San Jose State University
San Jose, California 95192 U.S.A.

Ho Kuen Ng

Department of Mathematics
San Jose State University
San Jose, California 95192 U.S.A.

Yihui Wen

Department of Mathematics
Suzhou Science and Technology College
Suzhou, Jiangsu 215009
People's Republic of China

Dedicated to Prof. M.K. Siu on the occasion of his 60th birthday

Abstract Let G be a (v, q) -graph. If the edges can be labeled $1, 2, 3, \dots, q$ so that the vertex sums are constant (mod v), then G is said to be edge-magic. A necessary condition of edge-magicness is $v \mid q(q+1)$. In [9], it was shown that for any graph G there is an integer k such that the k -fold graph $G[k]$ is edge-magic. The least such integer k is called the edge-magic index of G . We characterize k for some $(v, v+1)$ -graphs. If $v = p^n$, where p is a prime, then any (p^n, p^n+1) -graph has $p^n - 1$ as its edge-magic index. We also show here that there are infinitely many $(v, v+1)$ -graphs with edge-magic index less than $v - 1$.

1. Introduction. In this paper we consider graphs, possibly pseudographs and disconnected graphs without isolated vertices.

A graph $G = (V, E)$ with v vertices and q edges is said to be **magic** if there is a bijection $h : E \rightarrow \{1, 2, \dots, q\}$ such that the induced mapping $h^+ : V \rightarrow Z$, given by $h^+(u) = \sum \{h(u, w) : (u, w) \text{ in } E\}$ is a constant. It is said to be **edge-magic** if

there is a bijection $h : E \rightarrow \{1, 2, \dots, q\}$ such that the induced mapping $h^+ : V \rightarrow Z_v$, given by $h^+(u) = \sum\{h(u, w) : (u, w) \in E\} \pmod{v}$ is a constant.

The concept of edge-magic graphs was introduced by the first author, Seah and Tan [9]. A necessary condition for a (v, q) -graph to be edge-magic is $q(q+1) \equiv 0 \pmod{v}$. However, there are infinitely many connected graphs such as trees and cycles that satisfy the necessary condition but are not edge-magic.

The following figure shows three connected $(6, 9)$ -graphs that are edge-magic. (Figure 1)




Figure 1.

It was shown in [15] that the torus graph $C_m \times C_n$ is edge-magic for all $m, n > 2$.

In this paper we introduce a construction of edge-magic graphs by splitting edges. In [22], we introduced a general construction of multigraphs from a given simple graph. Let $N = \{1, 2, 3, \dots\}$ be the set of natural numbers. Given a pair (G, f) where $G = (V, E)$ is a simple graph and $f : E(G) \rightarrow N$, we can construct a graph $SPE(G, f)$ as follows: To each edge e of $E(G)$, if $f(e) = k$, we associate a set of parallel edges $P(e) = \{e_1, e_2, \dots, e_k\}$. We see that $V(SPE(G, f)) = V(G)$ and $E(SPE(G, f)) = \cup\{P(e) : e \in E(G)\}$.

We call the graph $SPE(G, f)$ the **edge-splitting extension graph** of (G, f) .

We illustrate here with an example:

Example 1. Let $G = C_4$ and $f : E(C_4) \rightarrow N$ be as follows:




Figure 2.

If $f: E(G) \rightarrow \mathbb{N}$ is a constant mapping, i.e., $f(e) = k$ for all edges e , we will denote $\text{SPE}(G, f)$ simply by $G[k]$. It was shown in [9] that for any graph G , there exists a k such that $G[k]$ is edge-magic. Denote by $\text{IM}(G)$ the set $\{k : G[k] \text{ is edge-magic}\}$. The smallest k in $\text{IM}(G)$ is called the **edge-magic index** of G .

Example 2. The following pseudograph of order 2 and size 3 has edge-magic index 1. Thus there exists a graph that is magic as well as edge-magic.




Figure 3.

The edge-magic indices of certain classes of graphs were considered in [16, 18, 19]. In this paper we determine the edge-magic indices of some $(v, v + 1)$ -graphs. Since vertex labels are defined using addition modulo v , the integers 0 and v are the same. They will be used interchangeably in this paper.

2. Edge-magic indices of $(v, v + 1)$ -graphs: Lower and upper bounds.

In this section we develop lower and upper bounds for the edge-magic index of a $(v, v + 1)$ -graph G .

Theorem 1. If G is a $(v, v + 1)$ -graph then its edge-magic index $\geq \min \{k : v \mid k(k + 1)\}$.

Proof. If m is the edge-magic index of G , then $G[m]$ is edge-magic. We observe that $q = (v + 1)m$. By the necessary condition of edge-magicness referred to above, we have $(v + 1)m[(v + 1)m + 1] \equiv 0 \pmod{v}$, giving $m^2 + m \equiv 0 \pmod{v}$. Thus $v \mid m(m + 1)$.

Let S be a set. We use $S \times n$ to denote the multi-set of n copies of S . Note that S may be a multi-set itself. From now on the term "set" means "multi-set" and set operations are multi-set operations.

Let S be a set of qk elements. A partition \wp of S is called a (q, k) -partition of S if each of the q classes in the partition contains k elements.

Notation. We let $[0] = \emptyset$, and for any integer $r > 0$, $[r] = \{1, 2, \dots, r\}$.

Suppose A is a set consisting of r integers, and its sum is s modulo v , we will say that A is an (s, r) -set or simply an s -set.

A mapping f is called a k -fold edge-magic labeling of a (v, q) -graph G if there is a (q, k) -partition \mathcal{P} of $[qk]$ such that $f : E(G) \rightarrow \mathcal{P}$ is a bijection and the induced mapping $f^+ : V(G) \rightarrow Z_v$ is a constant map, where

$$f^+(u) = \sum_{(u,w) \in E} \sum_{i \in f(u,w)} i \pmod{v}$$

Thus, finding an edge-magic labeling of $G[k]$ is equivalent to finding a k -fold edge-magic labeling of G .

The following example illustrates that $P_4[4]$ is edge-magic.

Example 3.

Consider the path P_4 and suppose $k = 4$. $[12] \equiv [4] \times 3 \pmod{4}$.




Figure 4.

Notation. For a cycle C_n with vertex set $\{x_1, x_2, \dots, x_n\}$, we denote by $C_n(t)$ the cycle with a chord (x_1, x_t) .

Example 4. The edge-magic index of $C_4(3)$ is 3. The 3-fold of $C_4(3)$ has 15 edges. We observe that $[15] \equiv [4] \times 3 \cup \{1, 2, 3\} \pmod{4}$. Figure 5 depicts a 3-fold edge-magic labeling of $C_4(3)$.




Figure 5.

We have the following general result.

Theorem 2. All $(4, 5)$ -graphs have edge-magic index 3.

Proof. By Theorem 1, 3 is the smallest possible value. Partition the integers from 1 to 15 (mod 4) into 5 sets, each with 3 integers, namely, $\{1, 1, 2\} \times 2$, $\{2, 3, 3\} \times 2$ and $\{0, 0, 0\}$. Each of these sets is a 0-set (mod 4), giving a 3-fold edge-magic labeling.

Note that the labeling in Theorem 2 is different from the one shown in Figure 5. In other words, there could be multiple edge-magic labelings.

Theorem 3. All $(5, 6)$ -graphs have edge-magic index 4.

Proof. Suppose G is a $(5, 6)$ -graph. By Theorem 1 we see that the edge-magic index is at least 4. Now $4 \times 6 = 24$ and $[24] = [4] \times 5 \cup \{0, 0, 0, 0\} \pmod{5}$.

Consider the partition \wp of $[24]$: $[4] \times 5, \{0, 0, 0, 0\}$. We see that each class is a 0-set. Therefore any bijection $f: E(G) \rightarrow \wp$ is a 4-fold edge-magic labeling.

These two Theorems will be generalized in Theorem 6 below.

Example 5. The following two $(5, 6)$ -graphs have edge-magic index 4.




Figure 6.

Remark. We want to note here that even if v divides $k(k+1)$, there is no guarantee that $G[k]$ is edge-magic. In other words, the edge-magic index could be strictly greater than such k . The following graph provides an example.




Figure 7.

Assume, on the contrary, that the edge-magic index = 2. Then there are 14 edges, with edge sum = $105 \equiv 3 \pmod{6}$. Let the sum of the two parallel edge labels be $a, b, c, d, e, f, g \pmod{6}$, as shown in the above diagram. Then $a + b + c + d + e + f + g \equiv 3 \pmod{6}$. Then, $g \equiv a + b + c + g \equiv a + d \equiv b + e \equiv c + f \equiv d + e + f \pmod{6}$. It follows that $a + b + c \equiv 0, a \equiv e + f, b \equiv d + f, c \equiv d + e$. Adding the last three congruences, we have $a + b + c \equiv 2(d + e + f)$. Since $g \equiv d$

$+ e + f \pmod{6}$ and $a + b + c \equiv 0 \pmod{6}$, we have $a + b + c + d + e + f + g \equiv 2(d + e + f) \equiv 0 \pmod{6}$, which is a contradiction. The minimum k such that 6 divides $k(k + 1)$ is 2. But $\min \text{IM}(G)$ is not 2.

We can show that the edge-magic index is not 3 by a similar argument. Assume not. Then $G[3]$, with 21 edges, is edge-magic. The total edge sum = $231 \equiv 3 \pmod{6}$. An argument identical to the above gives a contradiction.

Now we show that its edge-magic index = 5, by explicitly giving an edge-magic labeling (Figure 8). Note that this labeling will be discussed more generally in Theorem 5.





Figure 8.

Theorem 4. All connected (6, 7)-graphs, except the one depicted in Figure 8, have edge-magic indices 2.

Proof. Up to isomorphism, there are nineteen connected (6, 7)-graphs. We see that the edge-magic index is 2 for each of the following 18 graphs. (Figure 9)



For $v > 1$, partition the integers from 1 to $v^2 - 1 \pmod{v}$ into $v + 1$ sets, each with $v - 1$ integers, namely $[v - 1] \times v$, and $\{0, 0, \dots, 0\}$ (with $v - 1$ zeros). Each of these sets is a 0-set \pmod{v} , giving a $(v - 1)$ -fold magic labeling of G .

Case 2: v is even.

When $v = 2$, we have 2 vertices and 3 edges. $G[1] = G$ is obviously edge-magic.

For $v = 4$, see Theorem 2.

For $v = 6$, see Theorem 4 and the Remark before it.

Now consider $v \geq 8$. Partition the integers from 1 to $v^2 - 1 \pmod{v}$ into $v + 1$ sets, each with $v - 1$ integers, namely $\{0, 1, 2, \dots, (v/2) - 1, (v/2) + 1, \dots, v - 1\} \times (v - 1)$, $(\{v/2\} \times (v - 3)) \cup \{1, (v/2) - 1\}$, and $(\{v/2\} \times 3) \cup \{2, \dots, (v/2) - 2, (v/2) + 1, \dots, v - 1\}$. Each of the first $(v - 1)$ sets has a sum of $v(v - 1)/2 - (v/2) = v(v - 2)/2 \equiv 0 \pmod{v}$. The second to last set has a sum of $(v/2)(v - 3) + (v/2) = v(v - 2)/2 \equiv 0 \pmod{v}$. The last set has a sum of $3(v/2) + v(v - 1)/2 - 1 - ((v/2) - 1) - (v/2) = (v/2)v \equiv 0 \pmod{v}$. Thus each of these sets is a 0-set \pmod{v} , giving a $(v - 1)$ -fold edge-magic labeling of G .

3. $(v, v + 1)$ -graphs with $v = a$ prime power.

Theorem 6. Let p be a prime and n a positive integer. The edge-magic index of a $(p^n, p^n + 1)$ -graph is $p^n - 1$.

Proof. Consider two cases.

Case 1: $p = 2$.

Let the edge-magic index be k . The earlier necessary condition reduces to $2^n | k(k + 1)$. The smallest possible value of k satisfying this necessary condition is $2^n - 1$. It suffices to show that $G[2^n - 1]$ is edge-magic.

For $n = 1$, we have 2 vertices and 3 edges. $G[1] = G$ is obviously edge-magic.

For $n = 2$, see Theorem 2.

For $n \geq 3$, the proof of Theorem 5 gives such a labeling.

Case 2: p is odd.

Let the edge-magic index be k .

The earlier necessary condition reduces to $p^n | k(k + 1)$. The smallest possible value of k satisfying this necessary condition is $p^n - 1$. It suffices to show that $G[p^n - 1]$ is edge-magic. The proof of Theorem 5 gives such a labeling.

Corollary 7. All $(8, 9)$ -graphs have edge-magic indices 7.

Example 6. Consider $(8, 9)$ -graphs. The 0-sets $\pmod{8}$ are 7 copies of $\{0, 1, 2, 3, 5, 6, 7\}$, $\{4, 4, 4, 4, 4, 1, 3\}$, and $\{4, 4, 4, 2, 5, 6, 7\}$.

The edge-magic index of $C_8(5)$ is 7.




Figure 10.

Example 7. Consider $(16, 17)$ -graphs. We observe $16 = 2^4$, i.e., $n = 4$. The 0-sets (mod 16) are 15 copies of $\{0, 1, 2, \dots, 7, 9, \dots, 15\}$, $(\{8\} \times 13) \cup \{1, 7\}$, and $\{8, 8, 8, 2, 3, 4, 5, 6, 9, 10, 11, 12, 13, 14, 15\}$.

Example 8. The following $(7, 8)$ -graphs have edge-magic indices 6.




Figure 11.

We can consider all cases of $(v, v + 1)$ -graphs settled for $v \leq 11$. Note that $v = 2, 3, 4, 5, 7, 8, 9$, and 11 are primes or prime powers, and so are covered by Theorem 6. Theorem 4 deals with the case when $v = 6$. For $v = 10$, Theorem 1 shows that the edge-magic index ≥ 4 . Since $[44] = ([10] \times 4) \cup \{1, 2, 3, 4\}$, the following 0-sets show that the edge-magic index ≤ 4 : $\{0, 0, 0, 0\}$, $\{5, 5, 5, 5\}$, $\{1, 2, 3, 4\}$, two copies of $\{1, 1, 9, 9\}$, two copies of $\{2, 2, 8, 8\}$, two copies of $\{3, 3, 7, 7\}$, and two copies of $\{4, 4, 6, 6\}$.

4. Two families of $(v, v + 1)$ -graphs.

In this section we show that there exist infinitely many $(v, v + 1)$ -graphs with edge-magic index less than $v - 1$. We also show that for certain values of v , there can be different $(v, v + 1)$ -graphs with different edge-magic indices.

Construction 1:

Let $v = 2n(2n + 1)$, where n is a positive integer. Consider the graph $C_v(3)$, i.e., the cycle with v vertices, with an additional edge joining the first and third vertices. There are $v + 1$ edges, namely, the triangle connecting the first, second, and third vertices, and $2n(2n + 1) - 2$ other edges in the cycle C_v . We will show that it has edge-magic index $2n$.

Let the edge-magic index be k . The necessary condition earlier in this paper reduces to $2n(2n + 1) \mid k(k + 1)$. The smallest possible value of k satisfying this necessary condition is $2n$. It suffices to show that $C_v(3)[2n]$ is edge-magic.

In $C_v(3)[2n]$, there are $2n(2n(2n + 1) + 1)$ edges. Mod v , the set of edge labels = $(\{1, 2, \dots, 2n(2n + 1) - 1\} \leftrightarrow 2n) \uparrow (\{0\} \leftrightarrow 2n) \uparrow \{1, 2, \dots, 2n\}$. For the $2n(2n + 1) - 2$ edges other than the three forming the triangle, use the following 0-sets to label them: $\{1, 2n(2n + 1) - 1\} \leftrightarrow n$, $\{2, 2n(2n + 1) - 2\} \leftrightarrow n, \dots, \{n(2n + 1) - 1, n(2n + 1) + 1\} \leftrightarrow n$, with each of them used twice. Label the three edges in the triangle using $(\{0\} \leftrightarrow (2n - 1)) \uparrow \{n(2n + 1)\}$, $\{0\} \uparrow (\{n(2n + 1)\} \leftrightarrow (2n - 1))$, $\{1, 2, \dots, 2n\}$, each of which is an $n(2n + 1)$ -set (mod v). This gives a $(2n)$ -fold edge-magic labeling.

Example 9. Consider $n = 1$. We consider the $(6, 7)$ -graph $C_6(3)$. The 2-fold edge-magic labeling is given by the 3-sets $\{0, 3\}$, $\{0, 3\}$, $\{1, 2\}$ for the triangle, and the 0-sets $\{1, 5\}$, $\{1, 5\}$, $\{2, 4\}$, $\{2, 4\}$ for the other edges. Note that this is illustrated in the 15th case of Figure 9.

Example 10. Consider $n = 2$. We consider the $(20, 21)$ -graph $C_{20}(3)$. The 4-fold edge-magic labeling is given by the 10-sets $\{0, 0, 0, 10\}$, $\{0, 10, 10, 10\}$, $\{1, 2, 3, 4\}$ for the triangle, and the 0-sets $\{1, 1, 19, 19\}$, $\{1, 1, 19, 19\}$, $\{2, 2, 18, 18\}$, $\{2, 2, 18, 18\}$, \dots , $\{9, 9, 11, 11\}$, $\{9, 9, 11, 11\}$ for the other edges.

Construction 1 gives the following Theorem.

Theorem 8. There are infinitely many $(v, v + 1)$ -graphs with edge-magic index $< v - 1$.

Construction 2:

Consider the following $(v, v + 1)$ -graph, where $v = 2n(2n + 1)$, with n being an odd number. (Figure 12)




Figure 12.

We show that its edge-magic index is not $2n$. Assume the contrary, i.e., its edge-magic index is $2n$, and that the above diagram shows the sum of the parallel edge labels in $G[2n]$.

Mod $2n(2n+1)$, we have $h_1 \dots h_1 + h_2 \dots h_2 + h_3 \dots h_3 + h_4 \dots \dots h_{2n(2n+1)-7} + h_{2n(2n+1)-6} \dots h_{2n(2n+1)-6} + g \dots g + a + b + c \dots a + d \dots b + e \dots c + f \dots d + e + f$. Thus $h_2 \dots h_4 \dots \dots h_{2n(2n+1)-6} \dots 0, h_1 \dots h_3 \dots h_5 \dots \dots h_{2n(2n+1)-7} \dots g, a + b + c \dots 0, a \dots e + f, b \dots d + f, c \dots d + e$. From the last four congruences, we have $0 \dots a + b + c \dots 2(d + e + f)$. Since $g + a + b + c \dots d + e + f$, we have $g \dots d + e + f$. Thus the sum of all edge labels = $h_1 + h_2 + h_3 + h_4 + \dots + h_{2n(2n+1)-7} + h_{2n(2n+1)-6} + g + a + b + c + d + e + f \dots (n(2n+1) - 2)g + d + e + f \dots (n(2n+1) - 1)(d + e + f)$. Since n is odd, $n(2n+1) - 1$ is even, making the sum of all edge labels $\dots 0$.

On the other hand, the edge labels are $1, 2, \dots, 2n(2n(2n+1)+1)$. The sum is $(1 + 2n(2n(2n+1)+1))n(2n(2n+1)+1) \dots (1 + 2n)n$, which is not $0 \pmod{2n(2n+1)}$, giving a contradiction. Thus the edge-magic index cannot be $2n$.

Example 11. The graph in Figure 7 corresponds to the case $n = 1$ in this construction.

Constructions 1 and 2 give the following Theorem.

Theorem 9. There exist infinitely many values v such that for each such v , there exist two $(v, v+1)$ -graphs with different edge-magic indices.

5. Further work.

We might think that almost all $(v, v+1)$ -graphs have edge-magic index $v-1$, in the sense that if $A(v) = |\{(v, v+1)\text{-graphs having } v-1 \text{ as their edge-magic indices}|$ and $B(v) = |\text{all connected } (v, v+1)\text{-graphs}|$, then $\lim_{v \rightarrow \infty} A(v)/B(v) = 1$, as $v \rightarrow \infty$.

We will exhibit two classes of graphs to show that this conjecture is not true.

Consider $v = 2m$, where m is an odd number of the form $4k+1$, i.e., $v = 8k+2$. We show that all such $(v, v+1)$ graphs have edge-magic index $\leq m-1 < v-1$. Note that $[(m-1)(v+1)] = ([v] \times (m-1)) \cup [m-1]$, where $1+2+\dots+(m-1) = ((m-1)/2)m = 2km \equiv 0 \pmod{v}$. Label the $(2m+1)$ edges with the following $(0; m-1)$ -sets: $\{0, 0, \dots, 0\}$, $\{m, m, \dots, m\}$, $\{1, 2, \dots, m-1\}$, two copies of $(\{1\} \times ((m-1)/2)) \cup (\{v-1\} \times ((m-1)/2))$, two copies of $(\{2\} \times ((m-1)/2)) \cup (\{v-2\} \times ((m-1)/2))$, ..., and two copies of $(\{m-1\} \times ((m-1)/2)) \cup (\{m+1\} \times ((m-1)/2))$. Thus at least $1/8$ of the $(v, v+1)$ -graphs have edge-magic index $< v-1$.

Now consider $v = 2m$, where m is an odd number of the form $4k+3$, i.e., $v = 8k+6$. If the graph contains a triangle, we can modify our technique in Construction 1 to show that such graphs have edge-magic index $\leq m-1 < v-1$. Note that $[(m-1)(v+1)] = ([v] \times (m-1)) \cup [m-1]$, where $1+2+\dots+(m-1) = ((m-1)/2)m = (2k+1)m \equiv m \pmod{v}$. Label the 3 edges of the triangle using $(\{0\} \times (m-2)) \cup \{m\}$, $\{0\} \cup (\{m\} \times (m-2))$, and $\{1, 2, \dots, m-1\}$. Note that each of these is an $(m; m-1)$ -set. Label the other $(2m-2)$ edges using two copies of $(\{1\} \times ((m-1)/2)) \cup (\{v-1\} \times ((m-1)/2))$, two copies of $(\{2\} \times ((m-1)/2)) \cup (\{v-2\} \times ((m-1)/2))$, ..., and two copies of $(\{m-1\} \times ((m-1)/2)) \cup (\{m+1\} \times ((m-1)/2))$. Note that each of these is an $(0; m-1)$ -set. Thus these $(v, v+1)$ -graphs have edge-magic index $< v-1$.

We end this paper with a problem: What is the limit of $A(v)/B(v)$, as $v \rightarrow \infty$?

Acknowledgement. We express our thanks to the referee for his helpful suggestions.

References

- [1] J.A. Gallian, A dynamic survey of graph labeling, *The Electronic J. of Combin.*, (2001) # DS6, 1-144.
- [2] J. Keene and A. Simoson, Balanced strands for asymmetric, edge-graceful spiders, *Ars Combinatoria*, **42** (1996) 49-64.

- [3] L.M. Lee, S-M. Lee, and G. Murthy, On edge-graceful labeling of complete graphs – solutions of Lo's conjecture, *Congressum Numerantium*, **62** (1988) 225-233.
- [4] S-M. Lee, A conjecture on edge-graceful trees, *Scientia*, Ser. A, vol. **3** (1989) 45-57.
- [5] S-M. Lee, New directions in the theory of edge-graceful graphs, *Proceedings of the 6th Caribbean Conference on Combinatorics & Computing*, (1991) 216-231.
- [6] S-M. Lee, W.M. Pigg and T.J. Cox, on edge-magic cubic graphs conjecture, *Congressus Numerantium*, **105** (1994) 214-222.
- [7] S-M. Lee and E. Seah, Edge-graceful labelings of regular complete k-partite graphs, *Congressum Numerantium*, **75** (1990) 41-50.
- [8] S-M. Lee, E. Seah and S.P. Lo, On edge-graceful 2-regular graphs, *The Journal of Combinatoric Mathematics and Combinatoric Computing*, **12** (1992) 109-117.
- [9] S-M. Lee, E. Seah and S.K. Tan, On edge-magic graphs, *Congressus Numerantium*, **86** (1992) 179-191.
- [10] S-M. Lee, E. Seah, S-M. Tong, On the edge-magic and edge-graceful total graphs conjecture, *Congressus Numerantium*, **141** (1999) 37-48.
- [11] S-M. Lee, L. Wang, and Y.H. Wen, On edge-magic cubic multi graph and graphs, to appear in *Congressus Numerantium*.
- [12] S.P. Lo, On edge-graceful labelings of graphs, *Congressus Numerantium*, **50** (1985) 231-241.
- [13] J. Peng and W. Li, Edge-gracefulness of $C_m \times C_n$, *Proceedings of the Sixth Conference of Operations Research Society of China*, (Hong Kong: Global-Link Publishing Company), Changsha, October 10-15 (2000), 942-948.
- [14] K. Schaffer and S-M. Lee, Edge-graceful labelings of cartesian products, *Second China-U SA International Conference of Graph Theory, Combinatorics, Algorithms and Application*, San Francisco, 1989.
- [15] K. Schaffer and S-M. Lee, Edge-graceful and edge-magic labelings of Cartesian products of graphs, *Congressus Numerantium*, **141** (1999) 119-134.

- [16] W.C. Shiu, P.C.B. Lam and S-M. Lee, Edge-magicness of the composition of a cycle with a null graph, *Congressus Numerantium*, **132** (1998) 9-18.
- [17] W.C. Shiu, F.C.B. Lam and S-M. Lee, Edge-magic index sets of $(p, p - 1)$ -graphs, *Electronic Notes in Discrete Mathematics*, Vol. **11** (2002).
- [18] W.C. Shiu, F.C.B. Lam and S-M. Lee, Some edge-magic cubic graphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **40** (2002) 115-127.
- [19] W.C. Shiu, F.C.B. Lam and S-M. Lee, Edge-magic index sets of (p, p) -graphs, *Congressus Numerantium*, **137** (1999) 97-107.
- [20] W.C. Shiu, F.C.B. Lam and S-M. Lee, Edge-magic index sets of the second power of paths, preprint.
- [21] W.C. Shiu, S-M. Lee and K. Schaffer, Some k -fold edge-graceful labelings of $(p, p - 1)$ -graphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **38** (2001) 81-95.
- [22] Y.H. Wen, S-M. Lee and H. Sun, On edge-graceful splitting extension of graphs, manuscript.